



IMPA

NEWS

THE OFFICIAL NEWS LETTER OF THE INDEPENDENT MEDICAL PRACTITIONERS ASSOCIATION

IMPA News

- A very successful medical update programme on “Current Management of Dyslipidemia” by Dr. M.R. Mubarak (Consultant Cardiologist, Lanka Hospital, Colombo And “Non Alcoholic Fatty Liver Disease (NAFLD) / Non Alcoholic Steato Hepatitis (NASH)” by Dr.S.K.Kodisinghe, Consultant Gastroenterologist & Hepatologist, Teaching Hospital, Jaffna was held on Sunday 22nd July 2018 at the Lanka Hospital Auditorium sponsored by AstraZeneca (Hemas).
- CGP News
Pre congress - Sunday 16th September 2018 (2.00pm - 5.00pm) at the SLMA auditorium.
Inauguration - Saturday, 29th September 2018 at the Colombo Hilton Residencies.
Congress - Sunday, 30th September 2018 at the Colombo Hilton Residencies.
Post Congress - Sunday 7th October 2018 (2.00pm - 5.00pm) at the SLMA Auditorium.
- The Annual Sessions of the Vaccine Forum of Sri Lanka will be held on Sunday 7th October 2018 from 8.00 am - 1.00 pm at the New Lecture Theatre, Faculty of Medicine, Colombo.
- The IMPA council has approved the appointment of Mr. Ananda Kasthuriarachchi, Attorney at Law as the Honorable Legal Advisor of the IMPA by the initiative of Dr. H.L. Pathirajamudali (Hony. Treasurer of the IMPA)
- The IMPA council has approved the appointment of Dr (Mrs) Iyanthi Abeyewickrema as the IMPA representative in the AMR-NAP-TAC instead of Dr. Prasanna Siriwardana who wished to step down due to several other commitments.
- IMPA members are requested to forward articles to be included in the 2018 IMPA Journal which is due to be released at the AGM to be held in December 2018.
- The Primary Care Diabetic Group of Sri Lanka (PCDG) is organising a ‘DINISA’ (Free Diabetic Health Camp) on Sunday 28th October 2018 at the BMICH from 8.00 am - 4.00 pm.

Hema S Gopalan¹ • Anoop Misra^{1,2,3} • Ranil Jayawardena⁴

Received: 18 April 2018 / Revised: 9 May 2018 / Accepted: 11 May 2018

© Macmillan Publishers Limited, part of Springer Nature 2018

Introduction

Economic liberalization of India in the late 1990s ushered in an increase in life-expectancy, food security, urbanization, mechanization and shift towards a sedentary lifestyle and decreased energy expenditure. Together with an increased intake of energy-dense foods, consumption of alcohol and tobacco, these have resulted in nutritional imbalance, obesity and stress culminating into an ‘epidemic’ of type 2 diabetes mellitus (T2DM) and other non-communicable diseases (NCDs) [1]. To quantify deaths due to cardiovascular disease (CVD) and diabetes attributable to dietary and metabolic risks in South Asian countries, Yakoob and colleagues analysed data from the 2010 Global Burden of Disease national surveys and concluded that suboptimal diets were the leading cause. Such diets are characterised by low intake of fruits, whole grains, vegetables, and omega-3 polyunsaturated fatty acids [2]. In children and adolescents, nutritional risk factors associated with overweight / obesity include reduced fruit and vegetable consumption, consumption of energy-dense ‘junk’ foods and sugar-sweetened beverages (SSBs) [3].

In this review, we discuss determinants, emerging nutrition issues in social context, and some possible solutions for prevention of diabetes. The literature search has been carried using the terms ‘diet, nutrition or nutrient in Asian Indians or South Asians combined with ‘diabetes’ in the following

medical search databases; [1] PubMed (National Library of Medicine, Bethesda, MD) from 2006 to June, 2018, [2] Current Contents (Institute for Scientific Information, Thomson Scientific, Philadelphia, and [3] and Google Scholar. A manual search of the references was also carried out.

What is already known?**Intake of sugar and SSBs**

Intake of sugars from various sources is high in Asian Indians, owing to the traditional association of

offering of sugar-laden sweets in social and religious gatherings [4] (Fig. 1). SSBs, composed of energy-containing sweeteners are linked to T2DM, CVDs and obesity, even after adjustment for weight and total calorie consumption [5]. Though sales of SSBs have been static or declining in many high-income countries in recent years, the sales are rising in low- and middle-income countries (LMICs); India reports a 7% annual increase in sales [6]. Specifically, increased consumption of SSBs was observed in children and adolescents in a multi-country study and positively associated with excess body mass index [7].

Trans fats and food frying process (TFAs)

Adverse effects associated with excess consumption of trans-fatty acids (TFA) include dyslipidaemia, low-grade systemic inflammation and endothelial dysfunction, which may lead to CVDs [8] and affect insulin sensitivity possibly contributing to development of diabetes [9]. Oils conventionally utilized by South Asians contain high amount of TFAs (Vanaspati) or saturated fats [desi ghee (clarified butter), coconut oil]. Besides, the consumption of high fat snacks and fried street foods are prevalent in patients with T2DM in India [10]. Also, faulty cooking practices consisting of high-heat-treatment and frying in leftover oils, generates more TFAs regardless of the cooking oil used [11]. Regulations on utilization of low TFA oils and public awareness drives regarding unpropitious effects of frying/refrying must be done as a priority [12].

What we are getting to know now?**Processed foods**

Processed foods contain high amount of refined fats e.g. saturated fats and TFAs, sugar and salt that may contribute to obesity, T2DM, and CVDs. Between 1999 and 2017 intake of processed foods in India and Pakistan has increased 2-3 folds. In contrast, the developed Asian countries (Japan, Singapore) have reported either a decrease or a plateau in consumption of processed foods [13]. Further, though there is no


1. National-Diabetes, Obesity and Cholesterol Foundation (N-DOC), SDA, New Delhi 110016, India

2. Diabetes Foundation (India), New Delhi 110016, India

3. Fortis C-DOC Center of Excellence for Diabetes, Metabolic Diseases, and Endocrinology, B 16Chirag Enclave, New Delhi, India

4. Department of Physiology, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka

Fig. 1. Average intake of sugar globally vs. India (Reproduced with permission from Gulati and Misra [4])



national level data regarding sodium intake among South Asians; most of the South Asian snacks, curries and pickles contain high levels of salt [14] which may increase hypertension and CVD in south Asians with T2DM [15].

Vegetarian diets

South Asian countries like India and Nepal have a substantial population who are vegetarian, or those who, in addition, sparingly consume non-vegetarian foods. Although research on health effects of vegetarian diets is sparse, studies have shown improvement in insulin resistance in patients with T2DM consuming vegetarian diets [16]. However vegetarian diets of India often contain large amounts of carbohydrates and fried foods, which may have adverse metabolic consequences. Nevertheless, one study showed that vegetarian diets consumed by Asian Indians from India resulted in a better lipid profile and lower blood glucose levels than the non-vegetarians [17].

Need for nutrition guidelines

Traditionally, under-nutrition and nutritional deficiencies are the focus of nutrition researchers and policy makers in

South Asia. The existing nutritional guidelines rarely focus on NCDs, especially diabetes. Previously, we have highlighted the nutritional issues associated with diabetes among South Asians [18]. The currently available nutrition guidelines do not focus strongly on strategies for prevention and control of diabetes. Exercise guidelines for control and prevention obesity and diabetes are also not available for South Asian countries, except for India [19].

Erroneous beliefs and inadequate nutrition education in the population

Information transfer is becoming more and more rapid with the advent of mobile phones and computers in urban and in rural areas. Even though this has, to an extent, increased knowledge of nutrition, there is a possibility of people receiving incorrect messages or

‘half-truths’ [20]. Erroneous beliefs regarding the “health-value” of foods have been shown to influence consumption patterns in patients with diabetes in India [10]. Interestingly, in Sri Lanka, among all food and beverage-related advertisements on television, 78% were targeted at children and among these, 74% claimed health benefits without any disclaimers [21].

Non-compliance of medical nutrition therapy in type 2 diabetes

Diabetes-related self-care practices in South Asians are faulty with major non-compliance related to exercise and diets [22]. A study conducted in 11 cities of India showed that about 67% patients with T2DM faced major challenges in following correct diets [23]. Another study observed that about 19% of patients with diabetes were unaware of the role of diets in relation to glycaemic control [24]. Compliance to diets in South Asian patients with diabetes is affected by several factors; socio-economic strata, level of literacy, traditional dietary myths and practices, audio - visual media, and peer influences. Further, dietetic counselling services by trained personnel are available only with a small proportion of physicians. On most occasions, patients receive hasty

Table 1. Sources of Complex Carbohydrates

Type of food	English name	Hindi name
Cereals	Whole wheat	Gahun
	Brown rice	Chawal
Millets	Pearl Millet	Bajra
	Finger millet	Ragi
	Great millet /Sorghum	Jowar
Pulses	Red Gram	Tur Dal
	Green Gram	Sabut Moong
Legumes	Soya	Soya
	Peas	Muttar
	Horsegram	Kulthi

Source: Adapted from ref. [30]

Cont. on page 04

and incomplete dietary advice from their treating physicians [25]. Migrant South Asian patients state that cultural expectations and fear of alienation are reasons for the unrestricted food intake at social and community get-togethers where energy-dense fried foods and sweets are served [26]. It has also been demonstrated that improvement of dietary habits alone could help to reduce incident diabetes among pre-diabetic Asian Indian men [27].

Quality of carbohydrates

Carbohydrates in Indian diets in general and in patients with T2DM, are much higher than that recommended (64.1% of total dietary energy in one study) [28]. The focus however, should be to combine the following; decrease the 'quantity' and change 'quality' of the carbohydrate source, e.g., lesser-milled/high-quality/complex carbohydrates [29]. Table 1 shows primary sources of complex carbohydrates available in India [30]. In particular, millets are low-cost, low glycaemic index (GI), locally available complex carbohydrates, which are attractive alternative carbohydrate sources for patients with diabetes (Table 2).

Consumption of lesser processed grains; brown rice with legumes [41], pearl millet (Bajra) or barley instead of white bread [42], and wheat-based or millet-based formulations as dietary supplements instead of rice-based preparations [43] showed decreased glycaemic response among overweight, normoglycemic individuals and patients with T2DM. Millet-processed items for the modern consumers, e.g. leavened breads [44, 45], ready-to-cook flakes [46] and other products are also available along with efforts to include millets into staple diets (e.g., dosas or Indian pancakes or crepes) [38]. Overall, data regarding metabolic effects of complex carbohydrates in South Asians need further research.

What we need to know in the future?

Protein intake

Protein consumption among South Asians (particularly in India and Nepal) is low in quantity and quality, and is affected by factors like type of diet followed (vegetarianism) and the price-to-nutrient ratio. Pulses account for only 10 and 11% (rural and urban) of protein intake, while milk and dairy products contribute to 10 and 12% in rural and urban Indian diets, respectively. About 50–60% of protein intake in both rural and urban areas is from cereals [12, 47], the main source of protein intake in India.

Poor protein intake is associated with loss of muscle mass and function (Sarcopenia). Sarcopenia, as assessed by hand grip strength, is higher in South Asians as compared to Caucasians and Blacks and

is independently associated with T2DM [48]. To improve glucose-insulin metabolism in skeletal muscles, more protein intake along with a combination of aerobic and resistance physical activity is required in South Asians [49]. Further, to target obesity and poor muscle function at the same time, increasing protein in diet is a viable option in South Asians. Importantly, two recent studies in India have shown the beneficial effects of high-protein diets on multiple metabolic parameters including glycaemic control [12, 50].

Food choices

There could be several reasons and perceptions for staple food choices in South Asian populations. Factors responsible for dietary changes in rural regions of India are due to three primary causes (i) state-mandated fair price shops operated through the Public Distribution System (PDS) that sell refined wheat, sugar, palm oil, pulses, and kerosene fostering an environment to forego traditional staples in favour of products distributed by the PDS-nutrient-price ratio (ii) 'new' foods at low prices at local food stalls and shops (iii) shift in agricultural patterns due to the financial incentives of commercial crop production [29, 51].

A clinic-based study on high-income group patients with T2DM in an urban setting in north India revealed a paucity of knowledge level related to the healthy food choices and the basic concept of 'simple carbohydrates' and dietary fibre [10]. Another study in children and their mothers in north India showed incorrect notions about food (e.g., 'home-cooked meals are old-fashioned'); and mothers considered that feeding desi ghee to their children 'made the children strong' [52]. Another factor is the adoption of imbalanced urban-based diet and lifestyle by migrant populations from rural areas settled in urban slums [53]. These factors, along with epidemiological and nutrition transitions, need more studies.

Tools for dissemination of diet-related messages

Prevention/intervention programs and multi-level campaigns (audio-visual media, billboards and posters, street plays etc.) have been used to affect various health behaviours in India [54]. Further, e-health or m-health applications could be effectively utilized for prevention of T2DM [55, 56]. School-based educative interventions showed improvement in nutrition- and obesity-related knowledge, and anthropometric and biochemical profiles in children and adolescents [57–59]. Pandey et al. [60], with use of mass communication methods in a population-based intervention study, showed a significant increase in knowledge among women regarding importance of dietary proteins, fats, fibres and fruits.

Cont. on page 05

Table 2. Glycaemic Index of various Millets based food items as available in India

Type of millet	Food items prepared with ^a :- Local or hindi name of Millets	Glycaemic index (Mean value)	Indian states where commonly consumed
Major	Pearl millet	55.0 [31]	Gujarat and Rajasthan [32]
Major	Sorghum	44.32 [33]	Eastern and western Maharashtra, and northern Karnataka [34]
Minor	Finger millet	45.13 [35]	Karnataka, Tamil Nadu, Andhra Pradesh and parts of Uttarakhand [36]
Minor	Little Millet	52.54 [37]	Eastern Madhya Pradesh [38]
Minor	Foxtail millet	59.25 [39]	Karnataka, Tamil Nadu, Andhra Pradesh, Orissa, Bihar, Madhya Pradesh and Maharashtra
Minor	Barnyard millet	50.00 [40]	Uttar Pradesh, North-Eastern region and plains of North Bihar and Western Uttar Pradesh and Maharashtra

^a Food items may have more than one ingredient (other than millet)

It must be noted that for any individual behaviour change, cultural tailoring of health and nutrition messages should align with sociocultural belief systems with the use of multiple methods for information diffusion [61]. An apt example is a community-based participatory diabetes prevention

and management intervention in state of Gujarat, India, using community health workers that administered ten culturally and linguistically appropriate health education messages through face-to-face individual and group sessions (e.g. demonstrations of model meals and cooking techniques) that resulted in the significant improvement of knowledge regarding diabetes and CVDs, particularly in low socio-economic stratum [62]. Research on effective messages for prevention and control of diabetes tailored to geographical and cultural contexts in South Asia is clearly needed.

Food labelling

The Food Safety and Standards Authority of India (FSSAI) has introduced guidelines for advertising and labelling claims for packaged products providing information (nutritional value, weight and cost) regarding the ingredients. This regulation introduces transparency regarding high in fat, salt or sugar (HFSS) contents in processed foods. An expert group constituted by FSSAI for the food labelling of HFSS packaged/processed foods recommended ban on advertising of HFSS foods on children television channels and discouraged celebrity endorsements and recommended additional taxes on these (see below). Simple food labelling, which could be understood by all strata of society, its proper regulation, and its impact on T2DM and NCD is needed for all South Asian countries [63, 64].

Taxation

Basu et al. [65] suggested that a 20% excise tax on SSBs in India will prevent 11.2 million new cases of overweight and obesity (decrease of 3%), and 400,000 cases of T2DM (decrease of 1.6%) between 2014 and 2023. A similar tax (20%) on purchase of palm oil is estimated to result in a 1.3% reduction in cardiovascular deaths [66]. The state of Kerala in India has introduced a ‘fat tax’ (14.5%) that is added to HFSS foods sold at restaurants [67]. Impact of such taxation should be assessed, and if found to be effective, should be applied to all South Asian countries.

Neutraceuticals and natural compounds

A number of neutraceuticals and herbal compounds have been used for management of T2DM in South Asia, but few have been properly researched. There is emerging interest in cinnamon for improvement of metabolic parameters. In a recently conducted double-blind randomized trial on Asian Indians, use of cinnamon in individuals with metabolic syndrome lead to decrease in weight, glycaemic parameters, subclinical inflammation and hepatic transaminases [68]. Other locally available foods such as bitter

Cont. on page 06

gourd, curcumin, fenugreek seeds etc., need to be evaluated through well- designed clinical trials.

Conclusions

Epidemiological transition in South Asian countries has resulted in rural-urban migration, higher availability of processed foods (SSBs, fried foods), more purchasing power and a shift from traditional whole grain diets and high physical activity pattern to over-consumption of energy dense and poor physical activity. Coupled with inadequate health and nutrition-related information, these have become a recipe for an epidemic of NCDs particularly T2DM. Increasing awareness regarding importance of whole grain consumption, limiting sugar and saturated fat (fried foods) intake, use of healthier edible oils, and increase in protein intake through correctly positioned large scale nutrition education programs could help sensitize the community to adopt better health and nutrition practices. The government must strictly regulate HFSS foods in both unorganized and organized sectors and impose taxation on sales of SSBs and unhealthy oils. Research on nutraceuticals and functional foods requires further attention.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Misra A, Tandon N, Ebrahim S, Sattar N, Alam D, Shrivastava U, et al. Diabetes, cardiovascular disease, and chronic kidney disease in South Asia: current status and future directions. *BMJ*. 2017;11: j1420.
2. Yakoob MY, Micha R, Khatibzadeh S, Singh GM, Shi P, Ahsan H, et al. Impact of dietary and metabolic risk factors on cardiovascular and diabetes mortality in South Asia: analysis from the 2010 Global Burden of Disease Study. *Am J Public Health*. 2016;106:2113–25.
3. Jayawardena R, Ranasinghe P, Wijayabandara M, Hills AP, Misra A. Nutrition transition and obesity among teenagers and young adults in South Asia. *Curr Diabetes Rev*. 2017;13:444–51.
4. Gulati S, Misra A. Sugar intake, obesity, and diabetes in India. *Nutrients*. 2014;6:5955–74. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4277009/>
5. Malik VS, Popkin BM, Bray GA, Després J-P, Hu FB. Sugar sweetened beverages, obesity, type 2 diabetes and cardiovascular disease risk. *Circulation*. 2010;121:1356–64. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2862465/>.
6. WHO. Reducing consumption of sugar-sweetened beverages to reduce the risk of childhood overweight and obesity. WHO. http://www.who.int/elena/bbc/ssbs_childhood_obesity/en/. Accessed 28 Feb 2018.
7. Katzmarzyk PT, Broyles ST, Champagne CM, Chaput J-P, Fogelholm M, Hu G, et al. Relationship between soft drink consumption and obesity in 9–11 years old children in a multi-national study. *Nutrients*. 2016;8. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5188425/>.
8. Mozaffarian D, Katan MB, Ascherio A, Stampfer MJ, Willett WC. Trans fatty acids and cardiovascular disease. *N Engl J Med*. 2006;354:1601–13.
9. Guess N, Perreault L, Kerege A, Strauss A, Bergman BC. Dietary fatty acids differentially associate with fasting Versus 2-hour glucose homeostasis: implications for the management of subtypes of prediabetes. *PLOS ONE*. 2016;11:e0150148
10. Colles SL, Singh S, Kohli C, Mithal A. Dietary beliefs and eating patterns influence metabolic health in type 2 diabetes: a clinic-based study in urban North India. *Indian J Endocrinol Metab*. 2013;17:1066–72.
11. Bhardwaj S, Misra A, Gulati S, Anoop S, Kamal VK, Pandey RM, A randomized controlled trial to evaluate the effects of high protein complete (I A cto) VE geta R ian (PACER) diet in non-diabetics in North

- India. *Heliyon*. 2017;3:e00472. <http://linkinghub.elsevier.com/retrieve/pii/S2405844017305583>. Accessed 29 Mar 2018.
12. Gulati S, Misra A. Abdominal obesity and type 2 diabetes in Asian Indians: dietary strategies including edible oils, cooking practices and sugar intake. *Eur J Clin Nutrition*. 2017;71:850–7. <http://www.nature.com/doi/10.1038/ejcn.2017.92>. Cited 6 Apr 2018.
13. Baker P, Friel S. Processed foods and the nutrition transition: evidence from Asia. *Obes Rev*. 2014;15:564–77.
14. Mohan S, Prabhakaran D. Expert Meeting on Population Sodium Reduction Strategies for Prevention and Control of Non-communicable Diseases in the South-East Asia Region. 2012; http://www.searo.who.int/entity/noncommunicable_diseases/documents. Accessed 6 Apr 2018.
15. Devi P, Rao M, Sigamani A, Faruqui A, Jose M, Gupta R, et al. Prevalence, risk factors and awareness of hypertension in India: a systematic review. *J Hum Hypertens*. 2013;27:281–7.
16. Kahleova H, Matoulek M, Malinska H, Oliyarnik O, Kazdova L, Neskudla T, et al. Vegetarian diet improves insulin resistance and oxidative stress markers more than conventional diet in subjects with type 2 diabetes. *Diabet Med*. 2011;28:549–59.
17. Shridhar K, Dhillon PK, Bowen L, Kinra S, Bharathi AV, Prabhakaran D, et al. The Association between a vegetarian diet and cardiovascular disease (CVD) risk factors in India: the Indian migration study. *PLOS ONE*. 2014;9:e110586. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0110586>. 1272 H. S. Gopalan et al.
18. Misra A, Sharma R, Gulati S, Joshi SR, Sharma V, Ghafoorunissa G, et al. Consensus dietary guidelines for healthy living and prevention of obesity, the metabolic syndrome, diabetes, and related disorders in Asian Indians. *Diabetes Technol Ther*. 2011;13:683–94.
19. Misra A, Nigam P, Hills AP, Chadha DS, Sharma V, Deepak KK, et al. Consensus physical activity guidelines for Asian Indians. *Diabetes Technol Ther*. 2012;14:83–98.
20. Bureau AN. Can by consuming fruits one can cure diabetes? Viral Sach finds out. 2018. <http://www.abplive.in/india-news/can-by-consuming-fruits-one-can-cure-diabetes-viral-sach-finds-out-676596>. Accessed 2 Apr 2018.
21. Prathapan S, Wijewardena K, Low WY. Content analysis of food and beverages advertisements targeting children and adults on television in Sri Lanka. *Asia Pac J Public Health*. 2016;28(1 Suppl):86S–92S. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4785022/>. Accessed 16 Mar 2018.
22. Srinath KM, Basavegowda M, Tharuni NS. Diabetic self care practices in rural Mysuru, Southern Karnataka, India - a need for Diabetes Self Management Educational (DSME) program. *Diabetes Metab Syndr*. 2017;11(Suppl 1):S181–6.
23. Gudlavalleti MVS, Anchala R, Gudlavalleti ASV, Ramachandra SS, Shukla R, Jotheeswaran AT, et al. Perceptions and practices related to diabetes reported by persons with diabetes attending diabetic care clinics: The India 11-city 9-state study. *Indian J Endocrinol Metab*. 2016;20(Suppl 1):S26–32. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4847446/>. Accessed 26 Mar 2018.
24. Rai M, Kishore J. Myths about diabetes and its treatment in North Indian population. *Int J Diabetes Dev Ctries*. 2009;29:129–32. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2822217/>. Accessed 26 Mar 2018.
25. Patel M, Patel IM, Patel YM, Rathi SK. Factors associated with consumption of diabetic diet among type 2 diabetic subjects from Ahmedabad, Western India. *J Health Popul Nutr*. 2012;30:447–55. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3763616/>. Accessed 26 Mar 2018.
26. Lawton J, Ahmad N, Hanna L, Douglas M, Bains H, Hallowell N. ‘We should change ourselves, but we can’t’: accounts of food and eating practices amongst British Pakistanis and Indians with type 2 diabetes. *Ethn & Health*. 2008;13:305–19. <http://10.1080/13557850701882910>. Accessed 26 Mar 2018.
27. Ram J, Selvam S, Snehalatha C, Nanditha A, Simon M, Shetty AS, et al. Improvement in diet habits, independent of physical activity helps to reduce incident diabetes among prediabetic Asian Indian men. *Diabetes Res Clin Pract*. 2014;106:491–5.
28. Joshi SR, Bhansali A, Bajaj S, Bansal SS, Dharmalingam M, Gupta S, et al. Results from a dietary survey in an Indian T2DM population: a STARCH study. *BMJ Open*. 2014;4:e005138. <http://bmjopen.bmj.com/content/4/10/e005138>. Accessed 26 Mar 2018.
29. Mattei J, Malik V, Wedick NM, Hu FB, Spiegelman D, Willett WC, et al. Reducing the global burden of type 2 diabetes by improving the quality of staple foods: The Global Nutrition and Epidemiologic Transition Initiative. *Global Health*. 2015;11. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4489001/>.
30. Sarat Chandra K, Bansal M, Nair T, Iyengar SS, Gupta R, Manchanda SC, et al. Consensus statement on management of dyslipidemia in Indian subjects. *Indian Heart J*. 2014;66(Suppl 3): S1–51. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4297876/>.

Cont. on page 07

31. Mani UV, Prabhu BM, Damle SS, Mani I. Food & nutrition information. *Asia Pac J Clin Nutr.* 1993;2:111–4. <http://apjcn.nhri.org.tw/server/apjcn/2/3/111.htm>. Accessed 12 Mar 2018.
32. Basavaraj G, Rao PP, Bhagavatula S, Ahmed W. Availability and utilization of pearl millet in India. *SAT eJournal.* 2010;8:6. Available from: http://ejournal.icrisat.org/Volume8/IMPI/Availability_and_utilization.pdf.
33. Raidu V. Effect of processing and cooking on glycemic index of jowar varieties. *Int J Food Nutr Sci.* 2013;2:25–30.
34. Basavaraj G, Parthasarathy Rao P. Regional analysis of household consumption of Sorghum in Major Sorghum-producing and consuming states in India. Report No.: Working Paper Series no. 28. http://oar.icrisat.org/6422/1/J258_2011%20WPS%2028_full.pdf. Accessed 12 Mar 2018.
35. Shukla K, Srivastava S. Evaluation of finger millet incorporated noodles for nutritive value and glycemic index. *J Food Sci Technol.* 2014;51:527–34. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3931870/>. Accessed 12 Mar 2018.
36. Joseph SP. A Study on millets based cultivation and consumption in India. *Management Research.* 2013;2:10. Available from: <http://indianresearchjournals.com/pdf/IJMFMSR/2013/April/5.pdf>.
37. Patil KB, Chimmad BV, Itagi S. Glycemic index and quality evaluation of little millet (*Panicum miliare*) flakes with enhanced shelf life. *J Food Sci Technol.* 2015;52:6078–82. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4554602/>. Accessed 8 Mar 2018.
38. Narayanan J, Sanjeevi V, Rohini U, Trueman P, Viswanathan V. Postprandial glycaemic response of foxtail millet dosa in comparison to a rice dosa in patients with type 2 diabetes. *Indian J Med Res.* 2016;144:712–7. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5393082/>. Accessed 8 Mar 2018.
39. Nus Community. Kodo and kutki millets in Madhya Pradesh. <http://www.nuscommunity.org/research/projects/ifad-eu-ccafs-nus/kodo-and-kutki-millets-in-madhya-pradesh/>. Accessed 6 Apr 2018.
40. Ugare R, Chimmad B, Naik R, Bharati P, Itagi S. Glycemic index and significance of barnyard millet (*Echinochloa frumentacea*) in type II diabetics. *J Food Sci Technol.* 2014;51:392–5. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3907638/>. Accessed 8 Mar 2018.
41. Mohan V, Spiegelman D, Sudha V, Gayathri R, Hong B, Praseena K, et al. Effect of brown rice, white rice, and brown rice with legumes on blood glucose and insulin responses in overweight Asian Indians: a randomized controlled trial. *Diabetes Technol Ther.* 2014;16:317–25. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3996977/>. Accessed 9 Mar 2018.
42. Shukla K, Narain JP, Puri P, Gupta A, Bijlani RL, Mahapatra SC, et al. Glycaemic response to maize, bajra and barley. *Indian J Physiol Pharmacol.* 1991;35:249–54.
43. Shobana S, Kumari SRU, Malleshi NG, Ali SZ. Glycemic response of rice, wheat and finger millet based diabetic food formulations in normoglycemic subjects. *Int J Food Sci Nutr.* 2007;58:363–72. <https://doi.org/10.1080/09637480701252229>. Accessed 26 Mar 2018.
44. Chakraborty SK, Gupta S, Kotwaliwale N. Quality characteristics of gluten free bread from barnyard millet–soy flour blends. *J Food Sci Technol.* 2016;53:4308–15. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5223266/>.
45. Mannuramath M, Yenagi N, Orsat V. Quality evaluation of little millet (*Panicum miliare*) incorporated functional bread. *J Food Sci Technol.* 2015;52:8357–63. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4648869/>.
46. Takhellambam RD, Chimmad BV, Prkasam JN. Ready-to-cook millet flakes based on minor millets for modern consumer. *J Food Sci Technol.* 2016;53:1312–8. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4837707/>.
47. Kurpad AV, Minocha S. The health-nutrition-agriculture connect for protein in india. *Bull Nutr Found India.* 2017;38:8.
48. Ntuk UE, Celis-Morales CA, Mackay DF, Sattar N, Pell JP, Gill JMR. Association between grip strength and diabetes prevalence in black, South-Asian, and white European ethnic groups: a cross-Nutrition and diabetes in South Asia 1273 sectional analysis of 418 656 participants in the UK Biobank study. *Diabet Med.* 2017;34:1120–8.
49. Misra A, Alappan NK, Vikram NK, Goel K, Gupta N, Mittal K, et al. Effect of supervised progressive resistance-exercise training protocol on insulin sensitivity, glycemia, lipids, and body composition in asian indians with type 2 diabetes. *Diabetes Care.* 2008;31:1282–7. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2453659/>. Accessed 8 May 2018.
50. Gulati S, Misra A, Tiwari R, Sharma M, Pandey RM, Yadav CP. Effect of high-protein meal replacement on weight and cardio-metabolic profile in overweight/obese Asian Indians in North India. *Br J Nutr.* 2017;117:1531–40. https://www.cambridge.org/core/product/identifier/S0007114517001295/type/journal_article. Accessed 29 Mar 2018.
51. Little M, Humphries S, Patel K, Dewey C. Decoding the type 2 diabetes epidemic in rural India. *Med Anthropol.* 2017;36:96–110. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5359743/>.
52. Gulati S, Misra A, Colles SL, Kondal D, Gupta N, Goel K, et al. Dietary intakes and familial correlates of overweight/obesity: a four-cities study in India. *Ann Nutr Metab.* 2013;62:279–90.
53. Misra A, Pandey RM, Devi JR, Sharma R, Vikram NK, Khanna N. High prevalence of diabetes, obesity and dyslipidaemia in urban slum population in northern India. *Int J Obes Relat Metab Disord.* 2001;25:1722–9.
54. Daivadanam M, Absetz P, Sathish T, Thankappan KR, Fisher EB, Philip NE, et al. Lifestyle change in Kerala, India: needs assessment and planning for a community-based diabetes prevention trial. *BMC Public Health.* 2013;13:95. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3576354/>. Accessed 26 Mar 2018.
55. Ramachandran A, Snehalatha C, Ram J, Selvam S, Simon M, Nanditha A, et al. Effectiveness of mobile phone messaging in prevention of type 2 diabetes by lifestyle modification in men in India: a prospective, parallel-group, randomised controlled trial. *Lancet Diabetes & Endocrinol.* 2013;1:191–8. <http://linkinghub.elsevier.com/retrieve/pii/S2213858713700676>. Accessed 26 Mar 2018.
56. Muralidharan S, Mohan V, Anjana RM, Jena S, Tandon N, Allender S, et al. Mobile health technology (mDiab) for the prevention of type 2 diabetes: protocol for a randomized controlled trial. *JMIR Res Protoc.* 2017;6(12): e242. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5743924/>.
57. Shah P, Misra A, Gupta N, Kishore Hazra D, Gupta R, Seth P, et al. Improvement in nutrition-related knowledge and behaviour of urban Asian Indian school children: findings from the Medical education for children/adolescents for realistic prevention of obesity and diabetes and for healthy ageing (MARG) intervention study. *Br J Nutr.* 2010;104:427–36.
58. Panuganti P, Mehreen TS, Anjana RM, Mohan V, Mayer-Davis E, Ranjani H. Influence of socioeconomic status on knowledge of obesity and diabetes among adolescents in Chennai, South India. *Children (Basel).* 2017;4(8):61. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5575583/>. Accessed 19 Mar 2018.
59. Singhal N, Misra A, Shah P, Gulati S. Effects of controlled school-based multi-component model of nutrition and lifestyle interventions on behavior modification, anthropometry and metabolic risk profile of urban Asian Indian adolescents in North India. *Eur J Clin Nutr.* 2010;64:364–73.
60. Pandey RM, Agrawal A, Misra A, Vikram NK, Misra P, Dey S, et al. Population-based intervention for cardiovascular diseases related knowledge and behaviours in Asian Indian women. *Indian Heart J.* 2013;65:40–7. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3860769/>. Accessed 20 Mar 2018.
61. Mukherjee A, Underwood KC, Stewart AL, Ivey SL, Kanaya AM. Asian indian views on diet and health in the United States. *Fam Community Health.* 2013;36:311–23. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3987985/>.
62. Balagopal P, Kamamma N, Patel TG, Misra R. A community-based participatory diabetes prevention and management intervention in rural India using community health workers. *Diabetes Educ.* 2012;38:822–34.
63. Food Safety and Standards Authority of India (FSSAI). Report of Expert group on Consumption of Fat, Sugar, Salt & its Health effects on Indian population. FSSAI; 2017. https://www.fssai.gov.in/dam/jcr:4a9bc826.../Note_Report_HFSS_08_05_2017.pdf. Accessed 2 Mar 2017.
64. Food Labelling restrictions prescribed by FSSAI. Food Safety Helpline. 2015. <https://foodsafetyhelpline.com/2015/04/guidelines-on-fssai-restrictions-on-labels-of-food-products/cited>. Accessed 15 Mar 2018.
65. Basu S, Vellakkal S, Agrawal S, Stuckler D, Popkin B, Ebrahim S. Averting obesity and type 2 diabetes in india through sugar-sweetened beverage taxation: an economic-epidemiologic modeling study. *PLOS Med.* 2014;11:e1001582. <http://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1001582>. Accessed 28 Feb 2018.
66. Basu S, Babiarz KS, Ebrahim S, Vellakkal S, Stuckler D, Goldhaber-Fiebert JD. Palm oil taxes and cardiovascular disease mortality in India: economic-epidemiologic model. *BMJ.* 2013;347:f6048. <https://www.bmj.com/content/347/bmj.f6048>. Accessed 29 Mar 2018.
67. Indian State Imposes 'Fat Tax' on Fast Food. VOA. <https://leamingleish.voanews.com/a/southern-indian-state-of-kerala-imposes-fat-tax-on-fast-food/3426103.html>. Accessed 6 Mar 2018.
68. Gupta Jain S, Puri S, Misra A, Gulati S, Mani K. Effect of oral cinnamon intervention on metabolic profile and body composition of Asian Indians with metabolic syndrome: a randomized double-blind control trial. *Lipids Health Dis.* 2017;16:113. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5469078/>. Accessed 7 Mar 2018.

“Empowering the nation and upholding its cultural values...”
People’s Bank is a legacy that has been trusted for over 56 years.

Embarking on another milestone, we take pride in setting new trends in the banking and financial services industry, introducing fully digitized financial solutions to the nation.

Our revolutionary Self Banking Units have already begun to ease & improve our valuable customers’ lifestyles.

We stand steadfast as your dependable financial advisor, helping you nurture your life to new heights as you step in to the future.....

**STABILITY OF THE
FUTURE
DELIVERED TODAY...**



www.peoplesbank.lk

People's Bank is a licensed commercial bank supervised by the Central Bank of Sri Lanka.



**PEOPLE'S
BANK**

AA+ (Ika) Fitch Rating, AA Brand Finance Rating

PUBLISHED BY

INDEPENDENT MEDICAL PRACTITIONERS ASSOCIATION

275/75, PROF. STANLEY WIJESUNDARA MW, COLOMBO 7. Tel: 0112 501 113 Fax: 0112 500 818

E-mail: champa.impa@gmail.com | info@impa-lk.org Web: www.impa-lk.org